RESEALABLE BAG!

PRESS TO RESEAL

For outdoor use around the home only.

RESULTS Kills 40+ Types of Insects as listed GUARANTEED†

Spectracide®

TRIAZICIDE® INSECT KILLER

ONCE & DONE!®

GRANULES

Insecticida granulado

Season–Long Ant* Control

Kills on Contact	Once Treated Area has Been Watered and is Dry, Children and Pets Can Return to Lawn

BROADCAST WITH SPREADER™

*Excluding harvester and pharaoh ants.

For use on lawns or as a home barrier treatment

Active Ingredient:
Gamma-Cyhalothrin 0.05%
Other Ingredients 99.95%

†For terms of guarantee, see back panel.
‡Than standard 10,000 sq ft bags.

KEEP OUT OF REACH OF CHILDREN
CAUTION See back for additional precautionary statements.

MANTÉNGASE FUERA DEL ALCANCE DE LOS NIÑOS
PRECAUCIÓN Consulte la parte posterior para conocer las declaraciones preventivas adicionales.

Net Wt 20 lb (9.0 kg)

TREATS UP TO 25,000 SQ FT
2.5x MORE COVERAGE‡

RESULTS Kills 40+ Types of Insects as listed GUARANTEED†

Spectracide®

TRIAZICIDE® INSECT KILLER
ONCE & DONE!®
GRANULES
Insecticida granulado

TRIAZICIDE® INSECT KILLER

ONCE&DONE!®

GRANULES

Insecticida granulado

PRODUCT FACTS

WHAT IT DOES:
Helps protect your home from invading ants, except harvester and pharaoh ants. Also kills crickets, fire ants, chinch bugs, armyworms (fall and yellow-striped), billbugs, cutworms, earwigs, fleas, roaches (German and American), grubs, sod webworms, European crane flies, ticks and other listed insects.

WHERE TO USE:
Outdoors, on lawns and as a band treatment around house foundation.

QUESTIONS & COMMENTS:
Call 1-800-917-5438 or visit our website at **www.spectracide.com**

STOP **READ ENTIRE LABEL BEFORE USE**

DIRECTIONS FOR USE

It is a violation of Federal law to use this product in a manner inconsistent with its labeling.

STOP **Keep children and pets off treated area until product has been watered in and lawn is dry.**

In New York State, this product may not be applied to lawns within 100 ft of a coastal marsh, or stream that drains directly into a coastal marsh.

THREE EASY STEPS TO USE THIS PRODUCT

Step #1: Determine size of the job; measure the area to be treated.
Step #2: Use spreader setting chart to determine appropriate setting for your spreader.
Step #3: Treat as directed in rate chart. Thorough coverage is important. Water in granules immediately after application.

NOTE:
- Do not apply more than six (6) times per year per location at low use rate.
- Do not apply more than three (3) times per year per location at high use rate.
- Do not allow product to run off from target site or apply within 24 hours of rain.
- Do not make applications during rain. Apply this product directly to the lawn or garden area. Water treated area as directed on this label. Do not water to the point of runoff.
- Application is prohibited directly into sewers or drains, or to any area like a gutter where drainage to sewers, storm drains, water bodies or aquatic habitat can occur. Do not allow the product to enter any drain during or after application.

Treat the following pests during the months indicated. Make no more than six applications per year at low use rate or three times for highest use rate.

Jan	Feb	Mar	Apr	May	Jun	Jul	Aug	Sep	Oct	Nov	Dec
				Billbugs		Grubs					
		Dung Beetle Grubs									
			Hyperodes Weevils								
Treat ants (except harvester and pharaoh ants) and other insects when they first appear or you notice lawn damage.											

DIRECTIONS FOR APPLYING

Treatment Areas	Pests	How To Use	Use Rate
Band Treatment Around House Foundation	Ants*, fleas, roaches (German and American), palmetto bugs, water bugs, millipedes, mites, silverfish, sowbugs (pillbugs), crickets, earwigs	These pests are commonly found around or near foundations of houses as well as in lawns. Certain of these pests may enter houses. Apply a 3- to 5-foot band of granules next to the house foundation. Distribute granules uniformly. Water lightly immediately after application.	Apply at the rate of 0.8 lb per 1,000 sq ft (an area 5' x 200').
Overall Lawn Application	Ants*, armyworms (fall and yellow-striped), European crane flies, crickets, cutworms, earwigs, fleas, lawn moths (sod webworms), grasshoppers, Japanese beetle (adults), pillbugs, spittlebugs, mites, green June beetles, leafhoppers, millipedes, springtails and ticks (including ticks that may transmit Lyme Disease)	Apply with a fertilizer spreader. See spreader setting chart below. Apply when grass is dry. Water lawn lightly immediately after application. **For season-long control of fire ants, apply at higher use rate.**	Apply at the rate of 0.8 lb per 1,000 sq ft (an area 20' x 50'). **For season-long control of fire ants, apply at the rate of 2 lb per 1,000 sq ft (an area 20' x 50').**
	Billbugs, grubs (Japanese beetle grubs, green June beetle grubs), *Ataenius spretulus* (dung beetles), hyperodes weevils, mole crickets, chinch bugs	Immediately water grass thoroughly after application. *For billbug control,* apply when activity is first observed or when chewed or brown grass indicates damage from these insects. *For grub control,* except dung beetle, treat any time between late July and early October. *For dung beetle grub control,* apply in spring or early summer when damage is first noticed. Water grass thoroughly with ¼ to ½ inch of water immediately after application. Treat again in late summer or early fall if a second generation occurs.	Apply 1.2 lb per 1,000 sq ft of lawn.
Spot Treatment	Ants*	Water lightly immediately after application.	Apply ½ tsp over and around each anthill.

*Ants (except harvester and pharaoh): Argentine, carpenter, southern, field, Allegheny mound, cornfield, honey, pavement, red imported fire ant, nuisance

STORAGE AND DISPOSAL

Storage: Reclose container tightly after each use. Store in a cool, dry area which is inaccessible to children and pets.

Disposal: If empty: Nonrefillable container. Do not reuse or refill this container. Offer for recycling, if available.

If partly filled: Call your local solid waste agency for disposal instructions. Never place unused product down any indoor or outdoor drain.

PRECAUTIONARY STATEMENTS

Hazards to Humans and Domestic Animals

CAUTION. Causes moderate eye irritation. Avoid contact with eyes or clothing. Wash thoroughly with soap and water after handling and before eating, drinking, chewing gum or using tobacco. Food utensils such as spoons or measuring cups must not be used for food purposes after use in measuring pesticides.

FIRST AID

If in Eyes: Hold eye open and rinse slowly and gently with water for 15-20 minutes. Remove contact lenses, if present, after the first 5 minutes, then continue rinsing eye. Call a Poison Control Center or doctor for treatment advice.

Have the product container or label with you when calling a Poison Control Center or doctor, or going for treatment.

You may also contact 1-800-917-5438 for emergency medical treatment information.

Environmental Hazards

This product is highly toxic to fish and aquatic invertebrates. To protect the environment, do not allow pesticide to enter or run off into storm drains, drainage ditches, gutters or surface waters. Applying this product in calm weather when rain is not predicted for the next 24 hours will help to ensure that wind or rain does not blow or wash pesticide off the treatment area. Sweeping any product that lands on a driveway, sidewalk or street, back onto the treated area of the lawn or garden will help to prevent runoff to water bodies or drainage systems.

NOTICE: To the extent consistent with applicable law, buyer assumes all responsibility for safety and use not in accordance with directions.

Distributed by Spectrum Group
Division of United Industries Corporation
PO Box 142642, St. Louis, MO 63114-0642

EPA Reg. No. 9688-250-8845
EPA Est. Nos. 9198-OH-1(M), 9198-OH-2(B), 9198-AL-1(A), 8378-IN-1(CM), 69208-MO-1(AO), 39578-TX-1(AM), 62793-MO-1(AP), 58996-MO-1(AE), 5905-GA-1(CI), 5905-IA-1(CJ), 5905-AR-1(CK)
Circled letter is first letter of lot number. 11-11234 © 2010 UIC

Spreader Settings for Overall Lawn Application
Ajustes del difusor para la aplicación general en césped

Spreader Make and Model Marca y modelo del difusor	0.8 lb/1,000 sq ft 0.8 lb/1,000 pies cuadrados	1.2 lb/1,000 sq ft 1.2 lb/1,000 pies cuadrados	2 lb/1,000 sq ft 2 lb/1,000 pies cuadrados
CYCLONE			
Rotary	1	2	4
K-GRO			
Drop	2	3	5
REPUBLIC			
EZ Drop #530	3	5	8
EZ Rotary #540	1	1½	2½
SCOTTS			
AccuGreen 1000	3	3½	4½
AccuGreen 2000	3	3½	4½
AccuGreen 3000	3	3½	4½
SpeedyGreen 1000	2½	3½	4½
SpeedyGreen 2000	2½	3½	4½
SpeedyGreen 3000	2½	3½	4½
EdgeGuard	2½	3½	4½
EARTHWAY			
Rotary	5	6	7½
SEARS			
Drop	2½	3½	5
PRECISION			
Rotary (#S850)	1	2	3½
RED DEVIL			
Rotary	1	2	3½
TRUE TEMPER			
Rotary	1	1½	3

Note: The spreader settings listed are approximate. Accurately weigh a few pounds into your spreader and apply to a measured area. Then adjust spreader setting if necessary, in order to ensure correct rate of application.

†GUARANTEED RESULTS OR YOUR MONEY BACK

If you are not satisfied with this product for any reason, send your written request for a refund to the address at left, together with the original dated store receipt showing the price you paid. Please allow 6-8 weeks for processing.

ALTO **LEA TODA LA ETIQUETA ANTES DE USAR**

MODO DE EMPLEO

Se considera una violación a la ley federal usar este producto de una manera que no sea la indicada en la etiqueta.

ALTO **Mantenga los niños y las mascotas fuera de la zona tratada hasta que el producto se haya mojado y el césped esté seco.**

En el Estado de Nueva York, no se podrá aplicar este producto en céspedes ubicados a una distancia de hasta 100 pies de un pantano costero o de un arroyo que desemboque directamente en un pantano costero.

TRES PASOS FÁCILES PARA USAR ESTE PRODUCTO

Paso Nº 1: Determine la dimensión del trabajo; mida el área que se va a tratar.
Paso Nº 2: Use la tabla de ajuste del difusor para determinar el ajuste adecuado para el suyo.
Paso Nº 3: Efectúe el tratamiento como se indica en la tabla de dosis. Es importante que la cobertura sea completa. Moje los gránulos inmediatamente después de la aplicación.

NOTA:
- No aplique más de seis (6) veces por año por lugar a la dosis de uso más baja.
- No aplique más de tres (3) veces por año por lugar a la dosis de uso más alta.
- No permita que el producto se escurra del sitio que es el objetivo del tratamiento ni lo aplique hasta que hayan transcurrido 24 horas desde que llovió.
- No realice aplicaciones durante lluvias. Aplique este producto directamente al césped o zona del jardín. Riegue la zona tratada como se indica en esta etiqueta. No riegue hasta el punto de escurrimiento.
- Se prohíbe la aplicación directamente en alcantarillas o desagües, o en cualquier zona como una cuneta donde se pueda producir el desagüe hacia alcantarillas, alcantarillas pluviales, masas de aguas o hábitat acuático. No permita que el producto entre en ningún desagüe durante ni después de la aplicación.

Trate los siguientes insectos nocivos durante los meses indicados. No efectúe más de seis aplicaciones por año a la dosis de uso más baja o tres veces en el caso de dosis de uso más alta.

Ene	Feb	Mar	Abr	May	Jun	Jul	Ago	Sep	Oct	Nov	Dic
				Gorgojos		Larvas					
		Larvas del escarabajo del estiércol									
			Gorgojos (Hyperodes spp.)								
Trate las hormigas (excepto cosechera y faraona) y otros insectos cuando éstos aparezcan por primera vez o cuando observe daños en el césped.											

INSTRUCCIONES DE APLICACIÓN

Zonas de tratamiento	Insectos nocivos	Cómo usar	Dosis de uso
Tratamiento en franjas Alrededor de los cimientos de la casa	Hormigas*, pulgas, cucarachas (alemana y americana), cucarachas de caño, chinches de agua, milpiés, ácaros, lepismas, cochinillas de la humedad (cochinillas), grillos, tijeretas	Estos insectos nocivos se encuentran generalmente alrededor o cerca de los cimientos de las casas, así como también en los céspedes. Algunos de estos insectos nocivos pueden entrar en las casas. Aplique una franja de gránulos de 3 a 5 pies al lado de los cimientos de la casa. Distribuya los gránulos uniformemente. Riegue ligeramente inmediatamente después de la aplicación.	Aplique a la dosis de 0.8 libra por cada 1,000 pies cuadrados (un área de 5 pies x 200 pies).
Aplicación general en césped	Hormigas*, gusanos soldados (gusano cogollero y gusano pirero), típulas europeas, grillos, gusanos cortadores, tijeretas, pulgas, orugas del césped, saltamontes, escarabajos japoneses (adultos), cochinillas, cigarrillas, ácaros, escarabajos verdes de junio, saltarillas, milpiés, friesea y garrapatas (incluyendo las que pueden transmitir la enfermedad de Lyme)	Aplique con un difusor de fertilizante. Vea el cuadro de ajuste del difusor que se encuentra abajo. Aplique cuando el pasto esté seco. Riegue el césped ligeramente inmediatamente después de la aplicación. **Para el control estacional prolongado de las hormigas bravas, aplique la dosis de uso más alta.**	Aplique a la dosis de 0.8 libra por cada 1,000 pies cuadrados (un área de 20 pies x 50 pies). **Para el control estacional prolongado de las hormigas bravas aplique a la dosis de 2 libras por cada 1,000 pies cuadrados (un área de 20 pies x 50 pies).**
	Gorgojos, larvas (larvas de escarabajos japoneses, larvas de escarabajos verdes de junio), *Ataenius spretulus* (escarabajo del estiércol), gorgojos (Hyperodes spp.), grillos cebolleros, chinches	Moje completamente el pasto inmediatamente después de la aplicación. *Para el control de gorgojos,* aplique cuando comience a observar actividad o cuando el pasto masticado o marrón muestre el daño ocasionado por estos insectos. *Para el control de larvas,* excepto en el caso del escarabajo del estiércol, trate en cualquier momento durante el periodo que abarca desde fines de julio a principios de octubre. *Para el control de larvas de escarabajo del estiércol,* aplique en primavera o a principios del verano cuando se empieza a ver el daño. Moje completamente el pasto con ¼ a ½ pulgada de agua inmediatamente después de la aplicación. Si aparece una segunda generación, lleve a cabo un nuevo tratamiento a fines del verano o principios del otoño.	Aplique 1.2 libras por cada 1,000 pies cuadrados de césped.
Tratamiento localizado	Hormigas*	Riegue ligeramente inmediatamente después de la aplicación.	Aplique ½ cucharadita por encima y alrededor de cada hormiguero.

*Hormigas (excepto cosechera y faraona): Argentinas, carpinteras, del sur, de campo, Allegheny mound, de los campos de máiz, de la miel, del pavimento, hormigas bravas importadas, molestas

ALMACENAMIENTO Y ELIMINACIÓN

Almacenamiento: Vuelva a cerrar muy bien el recipiente después de cada uso. Guarde en un lugar fresco y seco que no sea accesible a los niños y a las mascotas.

Eliminación: Si está vacío: Envase no rellenable. No vuelva a usar ni rellenar este envase. Ofrézcalo para reciclar, si existe esa opción.

Si está parcialmente vacío: Llame a su servicio de eliminación de desechos más cercano para obtener instrucciones de eliminación. Nunca tire el producto sin usar por ningún drenaje interno ni externo.

DECLARACIONES PREVENTIVAS

Riesgos para los seres humanos y los animales domésticos

PRECAUCIÓN. Causa irritación moderada a los ojos. Evite el contacto con los ojos o la ropa. Lávese muy bien con agua y jabón después de manipularlo y antes de comer, beber, masticar chicle o usar tabaco. Los utensilios para alimentos como cucharas o tazas para medir no se deben usar con alimentos después de utilizarlos para medir pesticidas.

PRIMEROS AUXILIOS

Si entra en contacto con los ojos: Mantenga los ojos abiertos y enjuague con agua lentamente, con cuidado, durante 15 a 20 minutos. Si tiene lentes de contacto, retírelos después de los primeros 5 minutos, luego continúe enjuagando los ojos. Llame a un centro de control de envenenamientos o a un médico para recibir instrucciones de tratamiento.

Cuando llame a un centro de control de envenenamientos o a un médico, o intente obtener tratamiento, tenga a la mano el envase o la etiqueta del producto.

También puede llamar al 1-800-917-5438 para obtener información acerca del tratamiento médico de urgencia.

Riesgos para el medio ambiente

Este producto es extremadamente tóxico para los peces y los invertebrados acuáticos. Para proteger el medio ambiente, no permita que el pesticida entre o se escurra por alcantarillas pluviales, zanjas de drenaje, cunetas ni aguas superficiales. Aplicar este producto cuando haga buen tiempo cuando no se pronostique lluvia en las siguientes 24 horas, asegurará que ni el viento ni la lluvia sople o quite el pesticida de la zona de tratamiento. Barrer el producto que caiga en los caminos de entradas de automóviles, aceras o la calle y regresarlo a la zona de tratamiento del jardín o césped ayudará a evitar que escurra a masas de agua o sistemas de drenaje.

NOTA: Hasta el grado que sea compatible con la legislación pertinente, el comprador asume toda la responsabilidad para seguridad y uso que no esté de acuerdo con las instrucciones.

Nota: Los ajustes del difusor indicados son aproximados. Pese con exactitud algunas libras en su difusor y aplique a un área medida. Luego, si es necesario, ajuste el difusor, para asegurar la dosis de aplicación correcta.

PROOF OF PURCHASE
FPO UPC
0 71121 53960 1

RESEALABLE BAG!

RESULTS GUARANTEED†

Kills 40+ Types of Insects as listed

PRESS TO RESEAL

For outdoor use around the home only.

Spectracide®

TRIAZICIDE®
INSECT KILLER

ONCE&DONE!®

GRANULES

Insecticida granulado

Season–Long Ant* Control

Kills on Contact	Once Treated Area has Been Watered and is Dry, Children and Pets Can Return to Lawn

BROADCAST WITH SPREADER™

*Excluding harvester and pharaoh ants.

For use on lawns or as a home barrier treatment

Active Ingredient:
Gamma-Cyhalothrin 0.05%
Other Ingredients 99.95%

†For terms of guarantee, see back panel.
‡Than standard 10,000 sq ft bags.

KEEP OUT OF REACH OF CHILDREN
CAUTION See back for additional precautionary statements.

MANTÉNGASE FUERA DEL ALCANCE DE LOS NIÑOS
PRECAUCIÓN Consulte la parte posterior para conocer las declaraciones preventivas adicionales.

Net Wt 20 lb (9.0 kg)

TREATS UPTO **25,000** SQ FT
2.5x MORE COVERAGE‡

RESULTS Kills 40+ Types of Insects as listed GUARANTEED†

Spectracide®

TRIAZICIDE® INSECT KILLER
ONCE&DONE!®
GRANULES
Insecticida granulado

Spectracide®

TRIAZICIDE® INSECT KILLER

ONCE&DONE!®

GRANULES

Insecticida granulado

PRODUCT FACTS

WHAT IT DOES:
Helps protect your home from invading ants, except harvester and pharaoh ants. Also kills crickets, fire ants, chinch bugs, armyworms (fall and yellow-striped), billbugs, cutworms, earwigs, fleas, roaches (German and American), grubs, sod webworms, European crane flies, ticks and other listed insects.

WHERE TO USE:
Outdoors, on lawns and as a band treatment around house foundation.

QUESTIONS & COMMENTS:
Call 1-800-917-5438 or visit our website at **www.spectracide.com**

STOP **READ ENTIRE LABEL BEFORE USE**

DIRECTIONS FOR USE

It is a violation of Federal law to use this product in a manner inconsistent with its labeling.

STOP **Keep children and pets off treated area until product has been watered in and lawn is dry.**

In New York State, this product may not be applied to lawns within 100 ft of a coastal marsh, or stream that drains directly into a coastal marsh.

THREE EASY STEPS TO USE THIS PRODUCT

Step #1: Determine size of the job; measure the area to be treated.
Step #2: Use spreader setting chart to determine appropriate setting for your spreader.
Step #3: Treat as directed in rate chart. Thorough coverage is important. Water in granules immediately after application.

NOTE:
- Do not apply more than six (6) times per year per location at low use rate.
- Do not apply more than three (3) times per year per location at high use rate.
- Do not allow product to run off from target site or apply within 24 hours of rain.
- Do not make applications during rain. Apply this product directly to the lawn or garden area. Water treated area as directed on this label. Do not water to the point of runoff.
- Application is prohibited directly into sewers or drains, or to any area like a gutter where drainage to sewers, storm drains, water bodies or aquatic habitat can occur. Do not allow the product to enter any drain during or after application.

Treat the following pests during the months indicated. Make no more than six applications per year at low use rate or three times for highest use rate.

Jan	Feb	Mar	Apr	May	Jun	Jul	Aug	Sep	Oct	Nov	Dec
				Billbugs		Grubs					
		Dung Beetle Grubs									
			Hyperodes Weevils								
Treat ants (except harvester and pharaoh ants) and other insects when they first appear or you notice lawn damage.											

DIRECTIONS FOR APPLYING

Treatment Areas	Pests	How To Use	Use Rate
Band Treatment Around House Foundation	Ants*, fleas, roaches (German and American), palmetto bugs, water bugs, millipedes, mites, silverfish, sowbugs (pillbugs), crickets, earwigs	These pests are commonly found around or near foundations of houses as well as in lawns. Certain of these pests may enter houses. Apply a 3- to 5-foot band of granules next to the house foundation. Distribute granules uniformly. Water lightly immediately after application.	Apply at the rate of 0.8 lb per 1,000 sq ft (an area 5' x 200').
Overall Lawn Application	Ants*, armyworms (fall and yellow-striped), European crane	Apply with a fertilizer spreader. See spreader setting chart below. Apply when grass is dry. Water lawn	Apply at the rate of 0.8 lb per 1,000 sq ft

ALTO **LEA TODA LA ETIQUETA ANTES DE USAR**

MODO DE EMPLEO

Se considera una violación a la ley federal usar este producto de una manera que no sea la indicada en la etiqueta.

ALTO **Mantenga los niños y las mascotas fuera de la zona tratada hasta que el producto se haya mojado y el césped esté seco.**

En el Estado de Nueva York, no se podrá aplicar este producto en céspedes ubicados a una distancia de hasta 100 pies de un pantano costero o de un arroyo que desemboque directamente en un pantano costero.

TRES PASOS FÁCILES PARA USAR ESTE PRODUCTO

Paso Nº 1: Determine la dimensión del trabajo; mida el área que se va a tratar.
Paso Nº 2: Use la tabla de ajuste del difusor para determinar el ajuste adecuado para el suyo.
Paso Nº 3: Efectúe el tratamiento como se indica en la tabla de dosis. Es importante que la cobertura sea completa. Moje los gránulos inmediatamente después de la aplicación.

NOTA:
- No aplique más de seis (6) veces por año por lugar a la dosis de uso más baja.
- No aplique más de tres (3) veces por año por lugar a la dosis de uso más alta.
- No permita que el producto se escurra del sitio que es el objetivo del tratamiento ni lo aplique hasta que hayan transcurrido 24 horas desde que llovió.
- No realice aplicaciones durante lluvias. Aplique este producto directamente al césped o zona del jardín. Riegue la zona tratada como se indica en esta etiqueta. No riegue hasta el punto de escurrimiento.
- Se prohíbe la aplicación directamente en alcantarillas o desagües, o en cualquier zona como una cuneta donde se pueda producir el desagüe hacia alcantarillas, alcantarillas pluviales, masas de aguas o hábitat acuático. No permita que el producto entre en ningún desagüe durante ni después de la aplicación.

Trate los siguientes insectos nocivos durante los meses indicados. No efectúe más de seis aplicaciones por año a la dosis de uso más baja o tres veces en el caso de dosis de uso más alta.

Ene	Feb	Mar	Abr	May	Jun	Jul	Ago	Sep	Oct	Nov	Dic
				Gorgojos		Larvas					
		Larvas del escarabajo del estiércol									
			Gorgojos (Hyperodes spp.)								
Trate las hormigas (excepto cosechera y faraona) y otros insectos cuando éstos aparezcan por primera vez o cuando observe daños en el césped.											

INSTRUCCIONES DE APLICACIÓN

Zonas de tratamiento	Insectos nocivos	Cómo usar	Dosis de uso
Tratamiento en franjas Alrededor de los cimientos de la casa	Hormigas*, pulgas, cucarachas (alemana y americana), cucarachas de caño, chinches de agua, milpiés, ácaros, lepismas, cochinillas de la humedad (cochinillas), grillos, tijeretas	Estos insectos nocivos se encuentran generalmente alrededor o cerca de los cimientos de las casas, así como también en los céspedes. Algunos de estos insectos nocivos pueden entrar en las casas. Aplique una franja de gránulos de 3 a 5 pies al lado de los cimientos de la casa. Distribuya los gránulos uniformemente. Riegue ligeramente inmediatamente después de la aplicación.	Aplique a la dosis de 0.8 libra por cada 1,000 pies cuadrados (un área de 5 pies x 200 pies).

Overall Lawn Application	Ants*, armyworms (fall and yellow-striped), European crane flies, crickets, cutworms, earwigs, fleas, lawn moths (sod webworms), grasshoppers, Japanese beetle (adults), pillbugs, spittlebugs, mites, green June beetles, leafhoppers, millipedes, springtails and ticks (including ticks that may transmit Lyme Disease)	Apply with a fertilizer spreader. See spreader setting chart below. Apply when grass is dry. Water lawn lightly immediately after application. **For season-long control of fire ants, apply at higher use rate.**	Apply at the rate of 0.8 lb per 1,000 sq ft (an area 20' x 50'). **For season-long control of fire ants, apply at the rate of 2 lb per 1,000 sq ft (an area 20' x 50').**
	Billbugs, grubs (Japanese beetle grubs, green June beetle grubs), *Ataenius spretulus* (dung beetles), hyperodes weevils, mole crickets, chinch bugs	Immediately water grass thoroughly after application. *For billbug control,* apply when activity is first observed or when chewed or brown grass indicates damage from these insects. *For grub control,* except dung beetle, treat any time between late July and early October. *For dung beetle grub control,* apply in spring or early summer when damage is first noticed. Water grass thoroughly with ¼ to ½ inch of water immediately after application. Treat again in late summer or early fall if a second generation occurs.	Apply 1.2 lb per 1,000 sq ft of lawn.
Spot Treatment	Ants*	Water lightly immediately after application.	Apply ½ tsp over and around each anthill.

*Ants (except harvester and pharaoh): Argentine, carpenter, southern, field, Allegheny mound, cornfield, honey, pavement, red imported fire ant, nuisance

STORAGE AND DISPOSAL

Storage: Reclose container tightly after each use. Store in a cool, dry area which is inaccessible to children and pets.

Disposal: If empty: Nonrefillable container. Do not reuse or refill this container. Offer for recycling, if available.

If partly filled: Call your local solid waste agency for disposal instructions. Never place unused product down any indoor or outdoor drain.

PRECAUTIONARY STATEMENTS

Hazards to Humans and Domestic Animals

CAUTION. Causes moderate eye irritation. Avoid contact with eyes or clothing. Wash thoroughly with soap and water after handling and before eating, drinking, chewing gum or using tobacco. Food utensils such as spoons or measuring cups must not be used for food purposes after use in measuring pesticides.

FIRST AID

If in Eyes: Hold eye open and rinse slowly and gently with water for 15-20 minutes. Remove contact lenses, if present, after the first 5 minutes, then continue rinsing eye. Call a Poison Control Center or doctor for treatment advice.

Have the product container or label with you when calling a Poison Control Center or doctor, or going for treatment.

You may also contact 1-800-917-5438 for emergency medical treatment information.

Environmental Hazards

This product is highly toxic to fish and aquatic invertebrates. To protect the environment, do not allow pesticide to enter or run off into storm drains, drainage ditches, gutters or surface waters. Applying this product in calm weather when rain is not predicted for the next 24 hours will help to ensure that wind or rain does not blow or wash pesticide off the treatment area. Sweeping any product that lands on a driveway, sidewalk or street, back onto the treated area of the lawn or garden will help to prevent runoff to water bodies or drainage systems.

NOTICE: To the extent consistent with applicable law, buyer assumes all responsibility for safety and use not in accordance with directions.

Distributed by Spectrum Group
Division of United Industries Corporation
PO Box 142642, St. Louis, MO 63114-0642

EPA Reg. No. 9688-250-8845
EPA Est. Nos. 9198-OH-1(M), 9198-OH-2(B), 9198-AL-1(A), 8378-IN-1(CM), 69208-MO-1(AO), 39578-TX-1(AM), 62793-MO-1(AP), 58996-MO-1(AE), 5905-GA-1(CI), 5905-IA-1(CJ), 5905-AR-1(CK)
Circled letter is first letter of lot number. 11-11234 © 2010 UIC

Spreader Settings for Overall Lawn Application
Ajustes del difusor para la aplicación general en césped

Spreader Make and Model Marca y modelo del difusor	0.8 lb/1,000 sq ft 0.8 lb/1,000 pies cuadrados	1.2 lb/1,000 sq ft 1.2 lb/1,000 pies cuadrados	2 lb/1,000 sq ft 2 lb/1,000 pies cuadrados
CYCLONE			
Rotary	1	2	4
K-GRO			
Drop	2	3	5
REPUBLIC			
EZ Drop #530	3	5	8
EZ Rotary #540	1	1½	2½
SCOTTS			
AccuGreen 1000	3	3½	4½
AccuGreen 2000	3	3½	4½
AccuGreen 3000	3	3½	4½
SpeedyGreen 1000	2½	3½	4½
SpeedyGreen 2000	2½	3½	4½
SpeedyGreen 3000	2½	3½	4½
EdgeGuard	2½	3½	4½
EARTHWAY			
Rotary	5	6	7½
SEARS			
Drop	2½	3½	5
PRECISION			
Rotary (#S850)	1	2	3½
RED DEVIL			
Rotary	1	2	3½
TRUE TEMPER			
Rotary	1	1½	3

Note: The spreader settings listed are approximate. Accurately weigh a few pounds into your spreader and apply to a measured area. Then adjust spreader setting if necessary, in order to ensure correct rate of application.

†GUARANTEED RESULTS OR YOUR MONEY BACK

If you are not satisfied with this product for any reason, send your written request for a refund to the address at left, together with the original dated store receipt showing the price you paid. Please allow 6-8 weeks for processing.

Aplicación general en césped	Hormigas*, gusanos soldados (gusano cogollero y gusano pirero), típulas europeas, grillos, gusanos cortadores, tijeretas, pulgas, orugas del césped, saltamontes, escarabajos japoneses (adultos), cochinillas, cigarrillas, ácaros, escarabajos verdes de junio, saltarillas, milpiés, friesea y garrapatas (incluyendo las que pueden transmitir la enfermedad de Lyme)	Aplique con un difusor de fertilizante. Vea el cuadro de ajuste del difusor que se encuentra abajo. Aplique cuando el pasto esté seco. Riegue el césped ligeramente inmediatamente después de la aplicación. **Para el control estacional prolongado de las hormigas bravas, aplique la dosis de uso más alta.**	Aplique a la dosis de 0.8 libra por cada 1,000 pies cuadrados (un área de 20 pies x 50 pies). **Para el control estacional prolongado de las hormigas bravas aplique a la dosis de 2 libras por cada 1,000 pies cuadrados (un área de 20 pies x 50 pies).**
	Gorgojos, larvas (larvas de escarabajos japoneses, larvas de escarabajos verdes de junio), *Ataenius spretulus* (escarabajo del estiércol), gorgojos (Hyperodes spp.), grillos cebolleros, chinches	Moje completamente el pasto inmediatamente después de la aplicación. *Para el control de gorgojos,* aplique cuando comience a observar actividad o cuando el pasto masticado o marrón muestre el daño ocasionado por estos insectos. *Para el control de larvas,* excepto en el caso del escarabajo del estiércol, trate en cualquier momento durante el periodo que abarca desde fines de julio a principios de octubre. *Para el control de larvas de escarabajo del estiércol,* aplique en primavera o a principios del verano cuando se empieza a ver el daño. Moje completamente el pasto con ¼ a ½ pulgada de agua inmediatamente después de la aplicación. Si aparece una segunda generación, lleve a cabo un nuevo tratamiento a fines del verano o principios del otoño.	Aplique 1.2 libras por cada 1,000 pies cuadrados de césped.
Tratamiento localizado	Hormigas*	Riegue ligeramente inmediatamente después de la aplicación.	Aplique ½ cucharadita por encima y alrededor de cada hormiguero.

*Hormigas (excepto cosechera y faraona): Argentinas, carpinteras, del sur, de campo, Allegheny mound, de los campos de máiz, de la miel, del pavimento, hormigas bravas importadas, molestas

ALMACENAMIENTO Y ELIMINACIÓN

Almacenamiento: Vuelva a cerrar muy bien el recipiente después de cada uso. Guarde en un lugar fresco y seco que no sea accesible a los niños y a las mascotas.

Eliminación: Si está vacío: Envase no rellenable. No vuelva a usar ni rellenar este envase. Ofrézcalo para reciclar, si existe esa opción.

Si está parcialmente vacío: Llame a su servicio de eliminación de desechos más cercano para obtener instrucciones de eliminación. Nunca tire el producto sin usar por ningún drenaje interno ni externo.

DECLARACIONES PREVENTIVAS

Riesgos para los seres humanos y los animales domésticos

PRECAUCIÓN. Causa irritación moderada a los ojos. Evite el contacto con los ojos o la ropa. Lávese muy bien con agua y jabón después de manipularlo y antes de comer, beber, masticar chicle o usar tabaco. Los utensilios para alimentos como cucharas o tazas para medir no se deben usar con alimentos después de utilizarlos para medir pesticidas.

PRIMEROS AUXILIOS

Si entra en contacto con los ojos: Mantenga los ojos abiertos y enjuague con agua lentamente, con cuidado, durante 15 a 20 minutos. Si tiene lentes de contacto, retírelos después de los primeros 5 minutos, luego continúe enjuagando los ojos. Llame a un centro de control de envenenamientos o a un médico para recibir instrucciones de tratamiento.

Cuando llame a un centro de control de envenenamientos o a un médico, o intente obtener tratamiento, tenga a la mano el envase o la etiqueta del producto.

También puede llamar al 1-800-917-5438 para obtener información acerca del tratamiento médico de urgencia.

Riesgos para el medio ambiente

Este producto es extremadamente tóxico para los peces y los invertebrados acuáticos. Para proteger el medio ambiente, no permita que el pesticida entre o se escurra por alcantarillas pluviales, zanjas de drenaje, cunetas ni aguas superficiales. Aplicar este producto cuando haga buen tiempo cuando no se pronostique lluvia en las siguientes 24 horas, asegurará que ni el viento ni la lluvia sople o quite el pesticida de la zona de tratamiento. Barrer el producto que caiga en los caminos de entradas de automóviles, aceras o la calle y regresarlo a la zona de tratamiento del jardín o césped ayudará a evitar que escurra a masas de agua o sistemas de drenaje.

NOTA: Hasta el grado que sea compatible con la legislación pertinente, el comprador asume toda la responsabilidad para seguridad y uso que no esté de acuerdo con las instrucciones.

Nota: Los ajustes del difusor indicados son aproximados. Pese con exactitud algunas libras en su difusor y aplique a un área medida. Luego, si es necesario, ajuste el difusor, para asegurar la dosis de aplicación correcta.

PROOF OF PURCHASE
FPO UPC
0 71121 53960 1